



Research

# Effects of betaine and ascorbic acid on tonic immobility, superoxide dismutase and glutathione peroxidase in broiler chickens during the hot-dry season

Ifeanyichukwu Chukwuemeka Egbuniwe <sup>a</sup>  , Joseph Olusegun Ayo <sup>a</sup>, Mohammed Umar Kawu <sup>a</sup>, Aliyu Mohammed <sup>b</sup>

Show more 

 Share  Cite

<https://doi.org/10.1016/j.jveb.2015.11.001> 

[Get rights and content](#) 

## Abstract

The aim of the study was to evaluate effects of betaine and ascorbic acid (AA) on tonic immobility (TI), and activities of superoxide dismutase (SOD) and glutathione peroxidase (GPx) in broiler chickens, reared during the hot-dry season. Eighty White Ross breed day-old broilers were divided into 4 groups. Group I (control) was given sterile water; group II, betaine (250mg/kg); group III, AA (50mg/kg), and group IV, betaine+ AA, at 250+ 50mg/kg, respectively. The administration was done orally in drinking water and daily for 42 days. The duration of TI of each bird was measured. Serum obtained from blood samples collected from the birds were assayed for activities of the enzymes. The dry-bulb temperature (28.33°C–35.67°C), relative humidity (69.0–93.0%), and temperature-humidity index (27.85–36.1), recorded during the study period showed that the birds were subjected to heat stress. Betaine and its coadministration with AA decreased ( $P < 0.05$ ) TI duration compared to the control group. Betaine+ AA significantly increased ( $P < 0.01$ ) SOD activity compared with the controls. There were significant increases in GPx activity in broiler chickens in betaine ( $P < 0.01$ ), AA ( $P < 0.05$ ), and betaine+ AA ( $P < 0.001$ ) groups, when compared with that of controls. The result demonstrates that betaine and/or AA decreased TI and enhanced activities of the antioxidant enzymes in heat-stressed birds. It is concluded that the administration of betaine and/or AA to broiler chickens alleviates the adverse effects of heat stress by decreasing fearfulness and increasing activities of SOD and GPx enzymes.

## Introduction

High ambient temperature and high relative humidity (RH) occurring during the hot-dry season have been shown to cause heat stress in birds (Ayo et al., 2011, Minka and Ayo, 2012). Heat stress is a major environmental

stressor, posing a serious challenge to the success of poultry industry in the hot-humid regions of the world. The understanding of the biochemical changes in birds may be of value in controlling the negative effects caused by heat stress it causes (Lara and Rostagno, 2013). One component of stress in poultry is fear, and the duration of tonic immobility (TI) is the physiological indicator of fear. TI is less affected by external influences under the action of compulsory fixation. TI duration is also related to aggressive behavior, pecking, and cannibalism among hens in commercial poultry industry; hence, it is considered as an indirect diagnostic tool to assess stress, such as heat stress, in poultry (Tikhonov and Miftakhutdinov, 2014). TI is also an adaptive psychophysiological response (Sanotra et al., 2002). Studies have shown that genetic predispositions of poultry influence responses of birds to heat stress in terms of behavior and production (Mack et al., 2013). Birds confronted with treats exhibit fear-induced freezing called TI, mostly observed in prey species as a defense mechanism (Abe et al., 2013). Forkman et al. (2007) defined fear as an adaptive behavior which protects the animal from psychochemical damage, eliciting a reaction to the perception of actual danger. It is important to understand the behavior of animals because it affects their production and welfare (Broom, 2011). It is practically impossible to completely eliminate fear-inducing (stressful) situations during the rearing of birds (Lábaque et al., 2013), adversely affecting their welfare and production (de Haas et al., 2013). Zulkifli et al. (2009) described broiler chickens showing short or long TI responses as low-fear or high-fear responders, respectively. Antioxidant enzymes, including superoxide dismutase (SOD) and glutathione peroxidase (GPx), terminate free-radical reactions in cells by using a chain-reaction mechanism (Benzie, 2003). Enzymatic antioxidants like SOD and GPx scavenge both intracellular and extracellular oxidants, and prevent lipid peroxidation of cytomembranes (Sunil Kumar et al., 2011).

Betaine is widely found in nature, including animal tissues, seafood (from marine invertebrates), sugar beets, wheat bran, spinach, and microorganisms. It protects cells, protein, and enzymes from deleterious effects of environmental stress factors (like high ambient temperature, low water, high salinity) because it is an osmolyte (Craig, 2004). Betaine has been shown to enhance antioxidant defense, decreasing lipid peroxidation in the breast muscles of broiler chickens (Alirezai et al., 2012). Folate metabolism is adversely affected by insufficient serum levels of methyl donor such as betaine, resulting in decline in methylation and antioxidant capacity (Hamlin et al., 2013). It has been demonstrated that AA supplementation to heat-acclimated broiler chickens ameliorated the adverse effect of heat stress (El-Habbak et al., 2011). Ascorbic acid (AA) decreases the lipid oxidation in meat of broiler chickens stored for some days (Skřivan et al., 2012). It improves the breast meat yield in broilers during the hot-dry season (Abioja et al., 2011) and reduced the incidence of hypoxemia and susceptibility of broilers to heart failure (Nain et al., 2008). Glutathione plays a crucial role in controlling AA metabolism by regulating the expression and subcellular localization of the transporters, involved in the uptake of AA from extracellular sources (Mardones et al., 2012). Coadministration of antioxidants may be more effective than administration of single antioxidant because the combined administration has been shown to act synergistically to produce a magnified effect (Bai et al., 2014).

The aim of the study was to evaluate effects of coadministration of betaine and AA on TI, and the activities of SOD and GPx in broiler chickens during the hot-dry season.

---

## Section snippets

### Experimental site and thermal environment conditions

The experiment was carried out in a poultry house in Samaru—Zaria (11°10'N, 07°38'E), located in the Northern Guinea Savannah zone of Nigeria during the hot-dry season, from April to May, 2013 (Dzenda et al., 2013). ...

## Experimental birds and management

Eighty male broiler chickens, belonging to the White Ross breed, were purchased at day old from a commercial hatchery in Iperu-Remo (6°56'N, 3°43'E), Nigeria. Tags were tied to their legs for identification. The broiler chickens were maintained on commercial feed as follows: broiler ...

## Tonic immobility

The duration of TI in either the betaine-administered ( $77.87 \pm 6.62$ seconds) or betaine+ AA ( $62.71 \pm 4.22$ seconds) broiler chickens was shorter ( $P < 0.05$ ), when compared with that of the control birds ( $102.00 \pm 8.26$ seconds). The TI duration in betaine+AA-treated birds was also shorter ( $P < 0.05$ ) than that recorded in birds administered with AA alone ( $91.62 \pm 7.53$ seconds). Furthermore, TI duration in AA-treated birds was not different ( $P > 0.05$ ) from that of the control broilers ( $102.00 \pm ...$

## Tonic immobility and microenvironment

During the study period, the broiler chickens were reared under natural thermally stressful environmental conditions of the hot-dry season in the zone. The dry-bulb temperature (28°C-37°C), RH (68%-93%), and temperature-humidity index (27.85-36.10) values recorded during the study period fell predominantly outside the thermoneutral zone for broiler chickens. The normal ranges for dry-bulb temperature (18°C-21°C; Aengwanich and Simaraks, 2004), RH (65%-70%) (Akyuz and Boyain, 2010), and ...

## Acknowledgments

Every author cited in this article participated, at one point or the other, in its development. ...

---

## References (58)

H. Bai *et al.*

[Enhanced antioxidant effect of caffeic acid phenethyl ester and Trolox in combination against radiation induced-oxidative stress](#)

Chem. Biol. Interact (2014)

I.F. Benzie

[Evolution of dietary antioxidants](#)

Comp. Biochem. Physiol. (2003)

J.L. Campo *et al.*

[Realized heritability of tonic immobility in White-Leghorn hens: A replicated single generation test](#)

Poult. Sci. (1993)

S.A. Craig

[Betaine in human nutrition](#)

Am. J. Clin. Nutr. (2004)

E.N. de Haas *et al.*

## Fear, stress, and feather pecking in commercial white and brown laying hen parent-stock flocks and their relationships with production parameters

Poult. Sci. (2013)

B. Forkman *et al.*

## A critical review of fear test used on cattle, pigs, sheep, poultry and horses

Physiol. Behav. (2007)

X.H. Gu *et al.*

## Overexpression of heat shock protein 70 and its relationship to intestine under acute heat stress in broilers: 2. Intestinal oxidative stress

Poult. Sci. (2012)

M.C. Lábaque *et al.*

## Effects of thymol feed supplementation on female Japanese quail (*Coturnix coturnix*) behavioral fear response

Anim. Feed Sci. Technol. (2013)

H. Lin *et al.*

## Acute heat stress induces oxidative stress in broiler chickens

Comp. Biochem. Physiol. (2006)

B.D. Lott *et al.*

## Air velocity and high temperature effects on broiler performance

Poult. Sci. (1998)



View more references

---

## Cited by (23)

### Effects of Phenolic-Rich Onion (*Allium cepa* L.) Extract on the Growth Performance, Behavior, Intestinal Histology, Amino Acid Digestibility, Antioxidant Activity, and the Immune Status of Broiler Chickens ↗

2020, Frontiers in Veterinary Science

Show abstract ✓

### Functional roles of taurine, L-theanine, L-citrulline, and betaine during heat stress in poultry ↗

2022, Journal of Animal Science and Biotechnology

### Betaine: A Potential Nutritional Metabolite in the Poultry Industry ↗

2022, Animals

### Radical response: Effects of heat stress-induced oxidative stress on lipid metabolism in the avian liver ↗

2021, Antioxidants

## [Osmoregulatory function of betaine in alleviating heat stress in poultry ↗](#)

2018, Journal of Animal Physiology and Animal Nutrition

## [Betaine and antioxidants improve growth performance, breast muscle development and ameliorate thermoregulatory responses to cyclic heat exposure in broiler chickens ↗](#)

2018, Animals



[View all citing articles on Scopus ↗](#)

---

[View full text](#)

© 2015 Elsevier Inc. All rights reserved.



All content on this site: Copyright © 2025 Elsevier B.V., its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the relevant licensing terms apply.

